

# **Econ 102**

(Random walks  
and high finance)

Tom Carter

<http://astarte.csustan.edu/~tom/SFI-CSSS>

Fall, 2008

# Our general topics: ←

- ⊙ Some random (variable) background
- ⊙ What is a random walk?
- ⊙ Some Intuitive Derivations
- ⊙ Financial Modeling

# Some random (variable) background ←

This is a brief tour (peripatetic – wandering about) of some topics in random walks. It mostly consists of a few interesting ideas to start exploration of the general topic. We'll start with a bit on random variables.

- A random variable is a view into a set of possible values. Associated with each possible value is a probability. For a discrete random variable, there is a finite or countably infinite set of possible values and probabilities  $\{(a_i, p_i)\}$ , with the condition that  $\sum_i p_i = 1$ . Thus, for example, we could talk about the random variable  $\xi$  drawing values from the set of possible values  $\{(1, 1/2), (-1, 1/2)\}$ . When we evaluate the random variable  $\xi$ , we get either 1 or -1, each with probability 1/2.

- We can build new random variables. For example, if  $\xi_1$  and  $\xi_2$  are random variables over  $\{(1, 1/2), (-1, 1/2)\}$ , then  $\xi_1 + \xi_2$  is a random variable (but over the set  $\{(2, 1/4), (0, 1/2), (-2, 1/4)\}$ ). We must be a bit careful sometimes – for example,  $\xi_1 + \xi_1$  is also a random variable (but this time over  $\{(2, 1/2), (-2, 1/2)\}$ ).
- Given a random variable  $X$  over  $\{(a_i, p_i)\}$ , we define the *expectation* (or *expected value*) of  $X$  by:

$$\langle X \rangle = \sum_i p_i a_i.$$

Note that the expectation is a linear operator:

$$\langle \alpha X + \beta Y \rangle = \alpha \langle X \rangle + \beta \langle Y \rangle$$

for  $\alpha, \beta$  real numbers, and  $X, Y$  random variables.

- Note that the expectation of a constant  $\alpha$  is that constant (a constant can be thought of as a random variable over  $\{(\alpha, 1)\}$ ):

$$\langle \alpha \rangle = \alpha.$$

- Example: if  $\xi_1$  and  $\xi_2$  are random variables over  $\{(1, 1/2), (-1, 1/2)\}$ , then

$$\begin{aligned}\langle \xi_1 \rangle &= \frac{1}{2} * 1 + \frac{1}{2} * (-1) = 0 = \langle \xi_2 \rangle, \\ \langle \xi_1 + \xi_2 \rangle &= \langle \xi_1 \rangle + \langle \xi_2 \rangle = 0 + 0 = 0, \\ \langle \xi_1 \xi_2 \rangle &= \frac{1}{2} * 1 + \frac{1}{2} * (-1) = 0,\end{aligned}$$

but

$$\langle \xi_1^2 \rangle = \frac{1}{2} * 1^2 + \frac{1}{2} * (-1)^2 = 1,$$

and

$$\begin{aligned}\langle (\xi_1 + \xi_2)^2 \rangle &= \langle \xi_1^2 + 2 * \xi_1 \xi_2 + \xi_2^2 \rangle \\ &= \langle \xi_1^2 \rangle + 2 * \langle \xi_1 \xi_2 \rangle + \langle \xi_2^2 \rangle \\ &= 1 + 0 + 1 \\ &= 2.\end{aligned}$$

- Given a random variable  $X$ , we define the *variance* of  $X$  by:

$$V(X) = \langle (X - \langle X \rangle)^2 \rangle.$$

We can also calculate this as:

$$\begin{aligned} V(X) &= \langle (X - \langle X \rangle)^2 \rangle \\ &= \langle X^2 - 2X\langle X \rangle + \langle X \rangle^2 \rangle \\ &= \langle X^2 \rangle - 2\langle X\langle X \rangle \rangle + \langle \langle X \rangle^2 \rangle \\ &= \langle X^2 \rangle - 2\langle X \rangle^2 + \langle X \rangle^2 \\ &= \langle X^2 \rangle - \langle X \rangle^2. \end{aligned}$$

- The *standard deviation* of a random variable  $X$  is given by:

$$\sigma(X) = V(X)^{1/2}.$$

# What is a random walk? ←

There are a variety of ways to define a random walk. Here we start with a relatively simple version, which will allow us to develop some classical results on random walks. Later, we can generalize.

- Let  $\{\xi_i | i = 1, 2, 3, \dots\}$  be a set of (independent) random variables over  $\{(1, \frac{1}{2}), (-1, \frac{1}{2})\}$  (in particular, 'observing' one of the random variables has no effect on observations of any of the rest of them). Then a *simple random walk* is a sequence  $(S_n)$  where

$$S_0 = 0,$$

$$S_n = \xi_1 + \xi_2 + \dots + \xi_n.$$

- It is easy to see that

$$-n \leq S_n \leq n.$$

- We have that

$$\langle S_0 \rangle = 0,$$

and thus

$$\begin{aligned}\langle S_n \rangle &= \langle S_{n-1} + \xi_n \rangle \\ &= \langle S_{n-1} \rangle + \langle \xi_n \rangle \\ &= \langle S_{n-1} \rangle + 0 \\ &= \langle S_{n-1} \rangle \\ &= \langle S_{n-2} \rangle \\ &\vdots \\ &= \langle S_0 \rangle \\ &= 0.\end{aligned}$$

- We also have

$$\begin{aligned}
\langle S_n^2 \rangle &= \langle (S_{n-1} + \xi_n)^2 \rangle \\
&= \langle S_{n-1}^2 + 2 * S_{n-1} \xi_n + \xi_n^2 \rangle \\
&= \langle S_{n-1}^2 \rangle + 2 * \langle S_{n-1} \xi_n \rangle + \langle \xi_n^2 \rangle \\
&= \langle S_{n-1}^2 \rangle + 2 * \sum_{i=1}^{n-1} \langle \xi_i \xi_n \rangle + 1 \\
&= \langle S_{n-1}^2 \rangle + 2 * \sum_{i=1}^{n-1} 0 + 1 \\
&= \langle S_{n-1}^2 \rangle + 1 \\
&= \langle S_{n-2}^2 \rangle + 2 \\
&\vdots \\
&= \langle S_0^2 \rangle + n \\
&= n,
\end{aligned}$$

and thus

$$\begin{aligned}
V(S_n) &= \langle S_n^2 \rangle - \langle S_n \rangle^2 \\
&= n - 0 \\
&= n.
\end{aligned}$$

In other words, the variance of  $S_n$  is  $n$ , and the standard deviation of  $S_n$  is  $\sqrt{n}$ .

- We know that  $S_n$  ranges between  $-n$  and  $n$ . But how is it distributed across that range? In other words, if  $-n \leq k \leq n$ , what is the probability that  $S_n = k$  (i.e.,  $P(S_n = k)$ )? We can make a couple of observations. First, by symmetry,  $P(S_n = k) = P(S_n = -k)$ , since each of the  $\xi_i$  is over  $\{(1, \frac{1}{2}), (-1, \frac{1}{2})\}$ . Second, by a parity argument, if  $n$  is even and  $k$  is odd, or if  $n$  is odd and  $k$  is even, then  $P(S_n = k) = 0$ . Let us look, then, at the case  $n$  is even and  $k \geq 0$  is also even.

We know that  $S_n = \sum_i \xi_i$ , and that each of the  $\xi_i$  is either  $-1$  or  $1$ . Thus,  $S_n = k$  when exactly  $(\frac{n}{2} + \frac{k}{2})$  of the  $\xi_i$  are  $+1$  and the rest (i.e.,  $n - (\frac{n}{2} + \frac{k}{2}) = (\frac{n}{2} - \frac{k}{2})$ ) are  $-1$ . This can happen in  $\binom{n}{\frac{n}{2} + \frac{k}{2}}$  different ways. Each of these is equally likely, and there are  $2^n$  total possibilities.

Thus, we have that

$$\begin{aligned} P(S_n = k) &= \binom{n}{\frac{n}{2} + \frac{k}{2}} \frac{1}{2^n} \\ &= \frac{n!}{\left(\frac{n}{2} + \frac{k}{2}\right)! \left(n - \left(\frac{n}{2} + \frac{k}{2}\right)\right)! 2^n} \\ &= \frac{n!}{\left(\frac{n+k}{2}\right)! \left(\frac{n-k}{2}\right)! 2^n} \end{aligned}$$

# Some Intuitive Derivations



Every so often, I like to be a physicist (or biologist) and cavalier about error bounds.

- Suppose  $S_n = \sum_i \xi_i$  is a random walk, with  $\xi_i$  random variables over  $\{(1, \frac{1}{2}), (-1, \frac{1}{2})\}$ . Let us write  $P(S_n = k)$  as  $P(k, n)$ .

We can observe that

$$P(k, n + 1) = \frac{1}{2}P(k - 1, n) + \frac{1}{2}P(k + 1, n).$$

Now assume that  $n$  and  $k$  are large, let  $\delta$  and  $\tau$  be (small) real numbers, and then let  $x = \delta k$  and  $t = \tau n$ .

We then have  $P(x, t) = P(\delta k, \tau n)$ , and so

$$\begin{aligned} P(x, t + \tau) &= P(\delta k, \tau(n + 1)) \\ &= \frac{1}{2}P(\delta(k - 1), \tau n) + \frac{1}{2}P(\delta(k + 1), \tau n) \\ &= \frac{1}{2}P(x - \delta, t) + \frac{1}{2}P(x + \delta, t). \end{aligned}$$

From this, we get

$$\begin{aligned} P(x, t + \tau) - P(x, t) \\ = \frac{1}{2}(P(x - \delta, t) + P(x + \delta, t) - 2P(x, t)). \end{aligned}$$

Now consider two approximations. First, for small (infinitesimal)  $\tau$ , we have

$$P(x, t + \tau) = P(x, t) + \tau * \frac{\partial P(x, t)}{\partial t},$$

and for small (infinitesimal)  $\delta$ , we have

$$\begin{aligned} P(x + \delta, t) + P(x - \delta, t) \\ = 2P(x, t) + \delta^2 * \frac{\partial^2 P(x, t)}{\partial x^2}. \end{aligned}$$

Putting pieces together, we have:

$$\frac{\partial P(x, t)}{\partial t} = \frac{\delta^2}{2\tau} \frac{\partial^2 P(x, t)}{\partial x^2}$$

(i.e., the diffusion equation ...).

More generally, if we have a biased random walk (over  $\{(1, p), (-1, q)\}$ ), then  $P(x, t + \tau) = p * P(x - \delta, t) + q * P(x + \delta, t)$ , and, using the approximations, we have

$$\begin{aligned}
& \tau * \frac{\partial P(x, t)}{\partial t} \\
&= P(x, t + \tau) - P(x, t) \\
&= p * P(x - \delta, t) + q * P(x + \delta, t) - P(x, t) \\
&= p * \left( P(x, t) - \delta * \frac{\partial P(x, t)}{\partial x} + \frac{\delta^2}{2} * \frac{\partial^2 P(x, t)}{\partial x^2} \right) \\
&\quad + q * \left( P(x, t) + \delta * \frac{\partial P(x, t)}{\partial x} + \frac{\delta^2}{2} * \frac{\partial^2 P(x, t)}{\partial x^2} \right) \\
&\quad - P(x, t) \\
&= (p + q - 1)P(x, t) + (q - p) * \delta * \frac{\partial P(x, t)}{\partial x} \\
&\quad + \frac{(p + q)\delta^2}{2} * \frac{\partial^2 P(x, t)}{\partial x^2} \\
&= (q - p) * \delta * \frac{\partial P(x, t)}{\partial x} + \frac{\delta^2}{2} * \frac{\partial^2 P(x, t)}{\partial x^2}.
\end{aligned}$$

Writing this in a slightly different form,  
we have

$$\frac{\partial P(x, t)}{\partial t} = D * \frac{\partial^2 P(x, t)}{\partial x^2} + D * \beta * \frac{\partial P(x, t)}{\partial x},$$

where

$$D = \frac{\delta^2}{2\tau} \text{ and } \beta = \frac{2(1 - 2p)}{\delta}$$

(i.e., diffusion with drift ...).

- Let's look at another approach to continuous versions of these issues. Instead of looking at random variables over a discrete set, let the random variables draw their values from a probability distribution. In particular, let

$$w : \mathbb{R} \rightarrow [0, 1]$$

be an integrable (measurable) function, with

$$\int_{-\infty}^{\infty} w(s) ds = 1.$$

Then a random variable  $\xi$  over  $w(s)$  gives the value  $s$  with probability  $w(s)$  (i.e.,  $P(\xi = s) = w(s)$ ).

Note that we can use the Dirac delta function  $\delta(x - x_0)$  to recover the discrete examples if we want to.

Recall that the Dirac delta function has the properties

$$\delta(x - x_0)dx = 0 \quad \text{if } |x - x_0| > \frac{dx}{2}$$

$$\delta(x - x_0)dx = 1 \quad \text{if } |x - x_0| \leq \frac{dx}{2}$$

and

$$\int_{-\infty}^{\infty} \delta(x - x_0)dx = 1.$$

Then, if we let

$$w(x) = \sum_i p_i * \delta(x - a_i),$$

we are back in the discrete case.

Let's mention here also that the delta function has the property:

$$\delta(x - x_0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{it(x_0 - x)} dt.$$

Given a random variable  $\xi$  over a probability distribution  $w(s)$ , we can look at the expected value of the random variable

$$\langle \xi \rangle = \int_{-\infty}^{\infty} s * w(s) ds,$$

the mean square

$$\langle \xi^2 \rangle = \int_{-\infty}^{\infty} s^2 * w(s) ds,$$

the variance

$$V(\xi) = \langle \xi^2 \rangle - \langle \xi \rangle^2,$$

and so on.

Now suppose we have a probability distribution  $w(s)$  with mean  $\mu$  and standard deviation  $\sigma$  (i.e., if  $\xi$  is a random variable over  $w(s)$ , then  $\mu = \langle \xi \rangle$ , and  $\sigma^2 = V(\xi)$ ). Note that there is no guarantee for a given distribution  $w(s)$  that either the mean  $\mu$  or standard deviation  $\sigma$  exist – the integrals could diverge. In this example, we are assuming they do exist.

Let  $\{\xi_i\}$  be a collection of (independent) random variables over the distribution  $w(s)$ , and let  $S_n = S_0 + \sum_1^n \xi_i$  (with  $S_0 = 0$ ) be a random walk. What can we say about the distribution of the values of  $\frac{1}{n}S_n$ ? In other words, what can we say about the distribution of the average of  $n$  (identically distributed) random variables?

We want to find the probability that  $\frac{1}{n}S_n = x$  (let's write this as  $P_n(x)$ ). We will have  $\frac{1}{n}S_n = x$  if  $\xi_i = s_i$  and  $\frac{1}{n}\sum_i s_i = x$ . The probability of this happening is  $\prod_i w(s_i)$ , since the  $\xi_i$  are independent of each other. We need to add up the probabilities over all possible ways that  $\frac{1}{n}\sum_i s_i = x$  (as we did in the discrete case). In other words, we will have

$$P_n(x) = \int \int \cdots \int_{\frac{1}{n}\sum_i s_i = x} w(s_1) \cdots w(s_n) ds_1 \cdots ds_n.$$

The limits of integration are fairly messy, so we will use the Dirac delta function.

We will then have

$$P_n(x) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \delta\left(x - \frac{1}{n} \sum_j s_j\right) w(s_1) \cdots w(s_n) ds_1 \cdots ds_n.$$

Using the (fourier transform) property of the delta function, we have

$$\begin{aligned} 2\pi P_n(x) &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} e^{it\left(\frac{1}{n} \sum_j s_j - x\right)} w(s_1) \cdots w(s_n) dt ds_1 \cdots ds_n \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} e^{-itx} \prod_j \left( e^{it\frac{s_j}{n}} w(s_j) \right) dt ds_1 \cdots ds_n \\ &= \int_{-\infty}^{\infty} e^{-itx} \left( \prod_j \int_{-\infty}^{\infty} \left( e^{it\frac{s_j}{n}} w(s_j) \right) ds_j \right) dt. \end{aligned}$$

If we now let  $Q(t) = \int_{-\infty}^{\infty} e^{\frac{its}{n}} w(s) ds$ , we have

$$P_n(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} Q^n(t) dt.$$

Now let's look at  $Q(t)$ . We can expand the exponential to get

$$\begin{aligned}
 Q(t) &= \int_{-\infty}^{\infty} e^{\frac{its}{n}} w(s) ds \\
 &= \int_{-\infty}^{\infty} w(s) \left(1 + \frac{its}{n} - \frac{1}{2} \frac{t^2 s^2}{n^2} \dots\right) ds \\
 &= \int_{-\infty}^{\infty} w(s) ds + \frac{it}{n} \int_{-\infty}^{\infty} s * w(s) ds \\
 &\quad - \frac{t^2}{2n^2} \int_{-\infty}^{\infty} s^2 w(s) ds + \dots \\
 &= 1 + \frac{it}{n} \langle s \rangle - \frac{1}{2n^2} t^2 \langle s^2 \rangle + \dots
 \end{aligned}$$

Now we take the log, and use the expansion  $\ln(1 + y) = y - \frac{1}{2}y^2 + \dots$  to get

$$\begin{aligned}
 \ln(Q^n(t)) &= n \ln\left(1 + \frac{it}{n} \langle s \rangle - \frac{t^2}{2n^2} \langle s^2 \rangle + \dots\right) \\
 &= n * \left(\frac{it}{n} \langle s \rangle - \frac{1}{2n^2} t^2 \langle s^2 \rangle - \frac{1}{2} \left(\frac{it}{n} \langle s \rangle\right)^2 + \dots\right) \\
 &= \left(it \langle s \rangle - \frac{1}{2n} t^2 (\langle s^2 \rangle - \langle s \rangle^2) + \dots\right) \\
 &= \left(it \mu - \frac{1}{2n} t^2 \sigma^2 + \dots\right)
 \end{aligned}$$

where  $\mu$  and  $\sigma$  are the mean and standard deviation of the distribution  $w(s)$ .

Discarding all the higher order terms, and taking antilogs, we get

$$Q^n(t) = e^{it\mu - \frac{1}{2n}t^2\sigma^2},$$

and then that

$$P_n(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{it(\mu-x) - \frac{1}{2n}t^2\sigma^2} dt.$$

Now we use the formula

$$\int_{-\infty}^{\infty} e^{at-bt^2} dt = \sqrt{\frac{\pi}{b}} * e^{\left(\frac{a^2}{4b}\right)}$$

with  $a = i(\mu - x)$ ,  $b = \frac{1}{2n}\sigma^2$ , to finally get

$$\begin{aligned} P_n(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{it(\mu-x) - \frac{1}{2n}t^2\sigma^2} dt \\ &= \frac{1}{2\pi} \sqrt{\frac{\pi}{\frac{1}{2n}\sigma^2}} * e^{\left(-\frac{(\mu-x)^2}{4 * \frac{1}{2n}\sigma^2}\right)} \\ &= \frac{1}{\frac{\sigma}{\sqrt{n}} \sqrt{2\pi}} * e^{\left(-\frac{(x-\mu)^2}{2\left(\frac{\sigma^2}{n}\right)}\right)}. \end{aligned}$$

In other words, it is a normal distribution with mean  $\mu$  and standard deviation  $\frac{\sigma}{\sqrt{n}}$ .

Note that we made no assumptions about the distribution  $w(s)$  except that it actually has a mean ( $\mu$ ) and a standard deviation ( $\sigma$ ) (and, of course, that it goes to zero fast enough for large  $|s|$  that the approximations work out right . . . ).

What this says is that if we average a bunch of identically distributed independent random variables, the result is a normal distribution, whether or not the original distribution was normal.

This is usually called the *Central Limit Theorem*.

# Financial Modeling ←

Let's use some of these ideas to do some financial modeling. As an example, let's develop the (in)famous Black-Scholes model for options pricing.

- We can start with the simplest financial instrument, the fixed rate bond. If we “buy” amount  $V_0$  of a rate  $r$  bond at time  $t = 0$ , then at time  $t = 1$  we can redeem the for value  $V_0(1 + r)$ . If we wait longer to redeem the bond, then at some time in the future we can redeem the bond for

$$V(n, r) = V_0(1 + r)^n$$

One can also think of this as a “savings account” with interest rate  $r$ . In this case, we can ask the more general question, what is  $V(t, r)$  for real values of  $t$ , rather than just integral values of  $t$ ?

This will depend on the specifics of the bond (savings account). In its simplest form, the bond will have “coupons” that can be redeemed at specific times in the future, or in the case of a savings account, interest will be “compounded” on specific dates.

Let’s look at various possibilities for compounding. Suppose the bond has (annual) interest rate  $r$ . If interest is compounded  $k$  times during the year ( $k$  would be 4 for quarterly compounding, 12 for monthly compounding, etc.), then the value at time  $t$  would be

$$V(t, r, k) = V_0 \left(1 + \frac{r}{k}\right)^{kt}$$

If we smooth this out, and let  $k$  go to infinity (i.e., “continuous compounding”), then we will have

$$V(t, r) = V_0 \lim_{k \rightarrow \infty} \left(1 + \frac{r}{k}\right)^{kt} = V_0 e^{rt}.$$

We thus know how to set a price for a bond to be redeemed at some time  $t$  in the future.

- Now let's generalize. Suppose that the interest rate  $r$ , instead of being fixed, varied over time. What price should we be willing to pay for such a financial instrument? We can think of this financial instrument as a stock (share) in a corporation. Our "return on investment" will be uncertain, and will depend on the performance of the corporation (and also on the change in price of the stock). We have the potential to make a large profit (if the price of the stock goes up), but we now also have the potential to lose money (if the price of the stock goes down).

There is a difficulty here in that we don't really have the flexibility to buy the stock at whatever price we want today (depending on our calculation of the future value of the stock), but can only buy at today's price.

One thing we can do (at least potentially, assuming there are sellers willing), is to

purchase an “option” to buy the stock at some fixed price at some specific time in the future. Let’s simplify things a bit, and assume that over the time period in question, the stock will not pay any dividends (in other words, our profit/loss will only depend on changes in the price of the stock).

Once a market in “options” is developed, a variety of things become possible. Not only can we purchase options to buy a stock at a given price at a given time in the future (a “call” option), but we can also purchase an option to sell a stock at a given price at a given time in the future (a “put” option). Note that an option protects us against an adverse change in the price of the stock. For example, if we purchase a “call” option, and the price of the stock goes down, we let the option expire, and we lose only the amount we paid for the option (the premium). We

are thus protected against large losses. More generally, if we buy both “put” and “call” options, we can “hedge” our bets, and (we believe) protect ourselves against large losses, at the price of limiting the amount of gain we might make.

Our task, then, is to develop a model that will allow us to determine (estimate?) the price we should be willing to pay for an option.

In order to develop our model, we will have to make a variety of assumptions, many of them “simplifying” assumptions. It is possible (likely?) that at least some of our assumptions will be unrealistic, but at least will allow us to do computations. This presents us with an interesting dilemma – if our model is unrealistic, we may make very bad decisions if we depend on the model, but, if we make the model “realistic,” it may be useless to us because we can’t do the computations. Apparently, such is life . . .

- For this example, we're going to develop (a version of) the Black-Scholes option pricing model.

We'll have to make a variety of assumptions. These will include:

1. There is a completely safe (e.g., "FDIC insured savings account") fixed rate asset available.
2. There are "frictionless" markets (i.e., we can buy or sell any instrument at any time in any amount).
3. There are no transaction costs.
4. No "arbitrage" (there are no financial instruments that provide "risk free" profits above the fixed rate asset).

The most critical assumptions we will have to make concern the form of variability of the “price” of the stock on which we will be buying our options.

1. Variability is continuous (and possibly even smooth?). This will allow us to work in continuous time.
2. The distribution is “stable” (i.e., the distribution of the variability does not change over time).
3. Increments are independent (i.e., variability does not depend on history – there is no “memory” in the distribution).
4. The distribution has a finite mean and finite variance.
5. Variability is “independent of price” (i.e., the “value” of a change in price does not depend on the specific current price).

Putting all of this together, we will assume that the stock price  $x(t)$  gives us a continuous random variable

$$R(x, t) = \ln \left( \frac{x(t)}{x(0)} \right).$$

This random variable  $R(x, t)$  will be the return on the stock at time  $t$ , associated with stock price  $x(t)$ . More specifically, we will assume that the price  $x(t)$  satisfies the stochastic differential equation

$$dx(t) = \mu x(t) dt + \sigma x(t) dB_t$$

where  $B_t$  is Brownian motion,  $\mu$  is the “drift,” and  $\sigma$  is the volatility. This is generally called *geometric Brownian motion*. This equation has the solution

$$x(t) = x(0)e^{\left(\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma B_t\right)},$$

which is a log-normally distributed random variable with expected value  $\langle x(t) \rangle = x(0)e^{\mu t}$ , and variance  $\text{Var}(x(t)) = x(0)^2 e^{2\mu t} (e^{\sigma^2 t} - 1)$ .

The random variable  $R(x, t)$  is normally distributed, with mean  $(\mu - \sigma^2/2)t$  and variance  $\sigma^2 t$ .

We are interested in determining the current value of a financial instrument that will have a value in the future that depends on the price  $x(t)$  (or return  $R(x(t), t)$ ) of the underlying asset (stock). Let us call the value of this derived instrument  $w(x, t)$ . In the Black-Scholes case,  $w(x, t)$  will be the value of a “call” option.

To understand this better, we will study “portfolios” of various financial instruments. So, suppose we form a portfolio by putting one unit of our money into the secure interest bearing asset, and an amount  $-1/w_1$  (where  $w_1 = (\partial w / \partial x)$ ) in a “call” option with strike price  $K$  at time  $T$  (i.e., at time  $T$  we can buy the stock a price  $K$ ). Then the value of the portfolio will be  $p = x - w/w_1$ .

During a short period of time  $\Delta t$ , the value of the portfolio will change by  $\Delta p = \Delta x - \Delta w/w_1$ . We want to expand this formula, and fortunately there is a nice method (part of the *Ito calculus*) that allows us to write

$$\Delta w = \frac{\partial w}{\partial x} \Delta x + \frac{1}{2} \sigma^2 x^2 \frac{\partial^2 w}{\partial x \partial t} \Delta t + \frac{\partial w}{\partial t} \Delta t$$

(this is essentially the *chain rule* for *stochastic differentials*).

We then have

$$\Delta p = -\frac{1}{w_1} \left( \frac{1}{2} \sigma^2 x^2 w_{11} + w_2 \right) \Delta t$$

(where  $w_1, w_{11}$ , and  $w_2$  are the appropriate partial derivatives, and  $\sigma^2$  is a constant depending on  $w$ , essentially its variance or volatility).

Now we will use the “no arbitrage” assumption. Since  $\Delta p$  is an assured return, it must be that

$$\Delta p = r p \Delta t = r \left( x - \frac{w}{w_1} \right) \Delta t.$$

If we equate these two expressions for  $\Delta p$ , we get

$$-\frac{1}{w_1} \left( \frac{1}{2} \sigma^2 x^2 w_{11} + w_2 \right) \Delta t = r \left( x - \frac{w}{w_1} \right) \Delta t.$$

Dividing through by  $\Delta t$ , we can simplify to

$$w_2 + rxw_1 + \frac{1}{2} \sigma^2 x^2 w_{11} - rw = 0.$$

This is the Black-Scholes differential equation (B-S PDE) for  $w(x, t)$ . It has boundary conditions:

1.  $w(0, t) = 0$  for all  $t$ .
2.  $w(x, t) \sim x$  as  $t \rightarrow \infty$ .
3.  $w(x, T) = \max(x - K, 0)$  (recall  $K$  is the strike price at time  $T$ ).

- Our next task is to develop a solution to the B-S equation. We'll start by making some changes of variables. We'll reverse the order of time (with some normalization), since the “no arbitrage” rule allows us to use  $t = T$  for a boundary condition. We'll also move into a “logarithmic” mode (since the riskless asset grows exponentially), and normalize with respect to  $K$  (the strike price) – we shouldn't worry about the units of the price (e.g., dollars or pounds):

$$\begin{aligned}\tau &= \frac{\sigma^2}{2}(T - t) \\ z &= \ln(x/K) \\ v &= \frac{w}{K}\end{aligned}$$

Now we do some calculations:

$$\frac{\partial t}{\partial \tau} = -\frac{2}{\sigma^2}$$

and

$$\frac{\partial x}{\partial z} = x.$$

Then we have

$$\begin{aligned}\frac{\partial v}{\partial \tau} &= \frac{1}{K} \frac{\partial w}{\partial \tau} = \frac{1}{K} \left( \frac{\partial w}{\partial x} \frac{\partial x}{\partial \tau} + \frac{\partial w}{\partial t} \frac{\partial t}{\partial \tau} \right) \\ &= \frac{1}{K} \left( -\frac{2}{\sigma^2} \right) w_2\end{aligned}$$

and

$$\begin{aligned}\frac{\partial v}{\partial z} &= \frac{1}{K} \frac{\partial w}{\partial z} = \frac{1}{K} \left( \frac{\partial w}{\partial x} \frac{\partial x}{\partial z} + \frac{\partial w}{\partial t} \frac{\partial t}{\partial z} \right) \\ &= \frac{1}{K} x w_1.\end{aligned}$$

Using this, we get

$$\begin{aligned}\frac{\partial^2 v}{\partial z^2} &= \frac{1}{K} \frac{\partial x w_1}{\partial z} = \frac{1}{K} \left( \frac{\partial w_1}{\partial z} x + w_1 \frac{\partial x}{\partial z} \right) \\ &= \frac{1}{K} \left( \frac{\partial w_1}{\partial x} \frac{\partial x}{\partial z} x + \frac{\partial w_1}{\partial t} \frac{\partial t}{\partial z} x + x w_1 \right) \\ &= \frac{1}{K} (x^2 w_{11} + x w_1).\end{aligned}$$

Now we'll do some rearranging of the B-S equation, and then put in what we have gotten from the change of variables. Starting from

$$w_2 + rxw_1 + \frac{1}{2}\sigma^2 x^2 w_{11} - rw = 0.$$

we multiply by  $\frac{-2}{K\sigma^2}$

$$\frac{-2w_2}{K\sigma^2} + \frac{-2}{K\sigma^2}rxw_1 - \frac{1}{K}x^2 w_{11} + \frac{2}{K\sigma^2}rw = 0.$$

We rearrange, and add/subtract appropriate terms:

$$\begin{aligned} \frac{-2w_2}{K\sigma^2} &= \frac{x^2 w_{11}}{K} + \frac{2rxw_1}{K\sigma^2} - \frac{2rw}{K\sigma^2} \\ \frac{-2w_2}{K\sigma^2} &= \frac{x^2 w_{11}}{K} + \frac{w_1 x}{K} + \frac{2rxw_1}{K\sigma^2} + \frac{w_1 x}{K} - \frac{2rw}{K\sigma^2} \end{aligned}$$

or,

$$\begin{aligned} \frac{-2w_2}{K\sigma^2} &= \frac{1}{K}(x^2 w_{11} + xw_1) \\ &\quad + \left(\frac{2r}{\sigma^2} - 1\right) \frac{xw_1}{K} - \frac{2rw}{K\sigma^2} \end{aligned}$$

Then, putting in the change of variables, we have:

$$\frac{\partial v}{\partial \tau} = \frac{\partial^2 v}{\partial z^2} + (k - 1) \frac{\partial v}{\partial z} - kv$$

where  $k = \frac{2r}{\sigma^2}$ , and we have the boundary condition  $v(z, 0) = \max(e^z - 1, 0)$ .

This is close to a heat equation, but we need one more change of variables. If we let, for some constants  $\alpha, \beta$ ,

$$u(z, \tau) = e^{\alpha z + \beta \tau} v(z, \tau)$$

we will have

$$\begin{aligned} \beta u + \frac{\partial u}{\partial \tau} &= \alpha^2 u + 2\alpha \frac{\partial u}{\partial z} + \frac{\partial^2 u}{\partial z^2} \\ &\quad + (k - 1) \left( \alpha u + \frac{\partial u}{\partial z} \right) - ku \end{aligned}$$

If we choose

$$\alpha = -\frac{1}{2}(k - 1)$$

and

$$\beta = \alpha^2 + (k - 1)\alpha - k = -\frac{1}{4}(k + 1)^2,$$

then we are left with the heat equation:

$$\frac{\partial u}{\partial \tau} = \frac{\partial^2 u}{\partial z^2} .$$

At this point we'll just appeal to standard methods for solving the heat equation (e.g., Fourier series methods). The general solution will be of the form:

$$u(z, \tau) = \frac{1}{\sigma\sqrt{2\pi\tau}} \int_{-\infty}^{\infty} u_0(y) e^{-(z-y)^2/(2\sigma^2\tau)} dy.$$

Undoing the changes of variable, we get:

$$w(x, t; K, T, r) = x\Phi(d_1) - Ke^{-r(T-t)}\Phi(d_2) ,$$

where  $\Phi$  is the standard normal cumulative distribution function

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{(-\frac{u^2}{2})} du$$

and

$$d_1 = \frac{\ln(x/K) + (r + \frac{1}{2}\sigma^2)(T - t)}{\sigma\sqrt{T - t}}$$
$$d_2 = \frac{\ln(x/K) + (r - \frac{1}{2}\sigma^2)(T - t)}{\sigma\sqrt{T - t}}$$

càdlàg

[To top ←](#)

## References

- [1] Bachelier, Louis, *Theory of Speculation - The Origins of Modern Finance*, Translated and with Commentary by Mark Davis and Alison Etheridge, Princeton University Press, Princeton, 2006.
- [2] Spitzer, Frank, *Principles of Random Walk*, 2nd ed, Springer, New York, 1976.
- [3] Denny, Mark and Gaines, Steven, *Chance in Biology - Using Probability to Explore Nature*, Princeton University Press, Princeton, 2000.

[To top ←](#)